

**EMMA ID:** 02275

**Gene:** *CB2*

**Common name:** *Cnr2 tm2zim*

**Allele:** *Cnr2<sup>tm1Zim</sup>*

## Genotyping Information

Genotyping by end-point PCR based on gel is composed of a genespecific short range PCR using primers on wild type allele and a mutant allele-specific short range PCR. The combined results show the genotype of the mice. For example: mutant positive, wild type positive = Heterozygous.

### PCR primer pairs and expected size bands

| Assay    | Forward Primer | Reverse Primer | Expected Size Band (bp) |
|----------|----------------|----------------|-------------------------|
| Wildtype | mCB2           | CB2wt2         | 1100                    |
| Mutant   | mCB2           | pPNT           | 850                     |

### Primer sequences

| Primer Name | Sequence 5' --> 3'             |
|-------------|--------------------------------|
| mCB2        | AAATGCTTGATTGGTGTCAGCCTCTC     |
| pPNT        | TAAAGCGCATGCTCCAGACTGCCTT      |
| CB2wt2      | GGCTCCTAGGTGGTTTTTCACATCAGCCTC |

### PCR setup (Qiagen, Hot Start Plus)

| Component                 | Volume (µl) 1x | Final conc. |
|---------------------------|----------------|-------------|
| DNA (~ 50-100 ng)         | 2              |             |
| Q-Solution (5x)           | 2,5            | 0,5         |
| PCR-Buffer (10x)          | 2,5            | 1           |
| DNTP mix (10 mM)          | 0,5            | 0,2         |
| MgCl <sub>2</sub> (25 mM) | 1,5            | 1,5         |
| Primer 1 (10 pmol/µl)     | 1              | 0,4         |
| Primer 2 (10 pmol/µl)     | 1              | 0,4         |
| Taq Polymerase (5 U/µl)   | 0,3            | 0,06        |
| H <sub>2</sub> O*         | 13,7           |             |
| Final volume              | 25             |             |

\* The amount of H<sub>2</sub>O is adjusted with the number of primer.


### Amplification conditions

| PCR Settings                                       | Temperature (°C) | Time   | # of cycles |
|--|------------------|--------|-------------|
| 1 Denaturation<br>(Melting)                        | 95°C             | 5 min  | 1           |
| 2 Amplification<br>(Melting, Annealing,<br>Polym.) | 94°C             | 30 sec | 39          |
|  | 60°C             | 45 sec |             |
|  | 72°C             | 45 sec |             |
| 3 Polymerisation                                   | 72°C             | 10 min | 1           |
| 4 Cooling  | 12°C             | hold   | 1           |


These PCR conditions have been optimized for our methods and preparation kits. Adaptions may be required.

### Gel Image

WT-PCR



Mutant-PCR



Separated by gel electrophoresis on a 2% agarose gel.  
 ko/wt separate