

STRAIN NAME:**LEXKO 471****REAGENTS AND SOLUTIONS:**

- Taq polymerase 10X Buffer (1,5 mM MgCl₂) (MP QBiogene)
- dNTP (MPQ Biogene)
- Primers (Invitrogen)
- Taq polymerase (MPQBiogene)
- Mineral oil (EUROBIO)

REACTION MIX:

- Taq Buffer 1X
- dNTP (0,4 mM total)
- primer LEXKO 471 5' (0,24 µM)
- primer LEXKO 471 rev (0,48 µM)
- primer LTR2 (0,24 µM)
- Taq polymerase (40U/ml)
- DNA

PROGRAM:

5 min	94°C	
30 sec	94°C	
30 sec	60°C	35 cycles
1 min	72°C	
5 min	72°C	

MIGRATION :

Migration of the samples on a 1.5% agarose gel.

PRIMERS SEQUENCES

- Primer LEXKO 471 5' 5' AGCCTTGACCTTAAAGCGATTCACC 3'
- Primer LEXKO 471 rev 5' CTGCCTAGCTTAAGGACATG 3'
- Primer LTR2 5' AAATGGCGTTACTTAAGCTAGCTTGC 3'

AMPLIFICATION

Mutant allele : 174 bp
WT allele : 223 bp