

STRAIN NAME:**LEXKO 701****REAGENTS AND SOLUTIONS:**

- Taq polymerase 10X Buffer (1,5 mM MgCl₂) (MP QBiogene)
- dNTP (MPQ Biogene)
- Primers (Invitrogen)
- Taq polymerase (MPQBiogene)
- Mineral oil (EUROBIO)

REACTION MIX:

- Taq Buffer 1X
- dNTP (0,4 mM total)
- primer LEXKO 64 (0,48 µM)
- primer LEXKO 65 (0,24 µM)
- primer Neo3a (0,24 µM)
- Taq polymerase (40U/ml)
- DNA

PROGRAM:

5 min	94°C	
30 sec	94°C	
30 sec	60°C	35 cycles
1 min	72°C	
5 min	72°C	

MIGRATION :

Migration of the samples on a 1.5% agarose gel.

PRIMERS SEQUENCES

- Primer LEXKO 64 5' CTAACTGACCTTTAATCAGAGCAG 3'
- Primer LEXKO 65 5' CCATGTCTGCCCAAGTTCACTCTC 3'
- Primer Neo3a 5' GCAGCGCATCGCCTTCTATC 3'

AMPLIFICATION

Mutant allele : 311 bp
WT allele : 261 bp