

EMMA ID: 04125

Gene: *Fermt2*

Common name: *EPD0087_1_G04*

Allele: *Fermt2*^{tm1a(EUCOMM)Wtsi}

Allele Information

Further information about the allele can be found on IMPC website at (copy the link to web browser)
[http://www.mousephenotype.org/data/alleles/MGI:2385001/tm1a\(EUCOMM\)Wtsi](http://www.mousephenotype.org/data/alleles/MGI:2385001/tm1a(EUCOMM)Wtsi)

Links to the general information

About IKMC resource

<https://www.infrafrontier.eu/knowledgebase/protocols/ikmc-products>

IKMC allele types

<http://www.i-dcc.org/kb/entry/89/>

Allele conversion guide - genotyping tm1b, tm1c and tm1d mice (assays infos available when required)

<http://www.mousephenotype.org/about-ikmc/targeting-strategies>

IMPC mouse phenotype data, search by the gene name

<http://www.mousephenotype.org/>

Genotyping Information

Genotyping by end-point PCR based on gel is composed of a genespecific short range PCR using primers on wild type allele and a mutant allele-specific short range PCR. The combined results show the genotype of the mice.

For example: mutant positive, wild type positive = Heterozygous.

PCR primer pairs and expected size bands

| Assay | Forward Primer | Reverse Primer | Expected Size Band (bp) |
|----------|----------------|----------------|-------------------------|
| Mutant | Fermt2-5'arm | LAR3 | 450 |
| Wildtype | Fermt2-5'arm | Fermt2-3'arm | 743 |

Primer sequences

| Primer Name | Sequence 5' --> 3' |
|--------------|------------------------|
| Fermt2-5'arm | TACAGGTGGCTGACAAGATCC |
| Fermt2-3'arm | GTGAGGCTCACCTTCAGAGG |
| LAR3 | CAACGGGTTCTTCTGTTAGTCC |

PCR setup (Qiagen, Hot Start Plus)

| Component | Volume (μ l) 1x | Final conc. |
|-------------------------------|----------------------|-------------|
| DNA (~ 50-100 ng) | 2 | |
| Q-Solution (5x) | 2,5 | 0,5 |
| PCR-Buffer (10x) | 2,5 | 1 |
| DNTP mix (10 mM) | 0,5 | 0,2 |
| MgCl ₂ (25 mM) | 1,5 | 1,5 |
| Primer 1 (10 pmol/ μ l) | 1 | 0,4 |
| Primer 2 (10 pmol/ μ l) | 1 | 0,4 |
| Taq Polymerase (5 U/ μ l) | 0,3 | 0,06 |
| H ₂ O* | 13,7 | |
| Final volume | 25 | |


* The amount of H₂O is adjusted with the number of primer.

Amplification conditions

| PCR Settings | Temperature (°C) | Time | # of cycles |
|--|----------------------|----------------------------|-------------|
| 1 Denaturation (Melting) | 95°C | 5 min | 1 |
| 2 Amplification (Melting, Annealing, Polym.) | 94°C 65°C 72°C | 30 sec 45 sec 45 sec | 39 |
| 3 Polymerisation | 72°C | 10 min | 1 |
| 4 Cooling | 12°C | hold | 1 |

These PCR conditions have been optimized for our methods and preparation kits. Adoptions may be required.

Gel Image



Work as Triplex.

Separated by gel electrophoresis on a 2% agarose gel.

Genotyping using PCR-assays for cassette detection

LacZ reporter, Neo selection cassettes are inserted into the Knockout-first mutant allele. Cassette changes by allele conversion can be found on: <http://www.mousephenotype.org/about-ikmc/targeting-strategies>. For example, tm1b allele contains still lacZ reporter cassette, Neo selection cassette is deleted (promotor-driven only).

Please note that these assays are with universal cassette primers other than gene-specific. The confirmation on gene identity performed by e.g. sr genespecific PCR as provided is suggested .

PCR primer pairs and expected size bands

| Assay | Forward Primer | Reverse Primer | Expected Size Band (bp) |
|-------|--------------------|--------------------|----------------------------|
| lacZ | LacZ_multi_Deen_2F | LacZ_multi_Deen_2R | mut 81 bp,wt without band |
| Neo | Neo_long_Deen_F1 | Neo_long_Deen_R1 | mut 186 bp,wt without band |

Primer sequences

| Primer Name | Sequence 5' --> 3' |
|--------------------|------------------------|
| LacZ_multi_Deen_2F | TACTGGAGGCTGAAGTTCAGAT |
| LacZ_multi_Deen_2R | GCGTTTCACCCCTGCCATAA |
| Neo_long_Deen_F1 | TTGAACAAGATGGATTGCACGC |
| Neo_long_Deen_R1 | CCTCGTCCTGCAGTTCATT |

PCR setup (Qiagen, Hot Start Plus)

Amplification conditions

| Component | Volume (µl) | Final conc. | PCR Settings | Temperature (°C) | Time | # of cycles |
|-------------------------|-------------|-------------|--|----------------------|----------------------------|-------------|
| DNA (~ 50-100 ng) | 2 | | Denaturation (Melting) | 95°C | 5 min | 1 |
| Q-Solution (5x) | 2,5 | 0,5 | Amplification (Melting, An- nealing, Polym.) | 94°C 58°C 72°C | 30 sec 45 sec 45 sec | 39 |
| PCR-Buffer (10x) | 2,5 | 1 | | | | |
| DNTP mix (10 mM) | 0,5 | 0,2 | | | | |
| MgCl2 (25mM) | 1,5 | 1,5 | Polymerisation | 72°C | 10 min | 1 |
| Primer 1 (10 pmol/µl) | 1 | 0,4 | | | | |
| Primer 2 (10 pmol/µl) | 1 | 0,4 | | | | |
| Taq Polymerase (5 U/µl) | 0,3 | 0,06 | | | | |
| H2O | 13,7 | | Cooling | 12°C | hold | 1 |
| Final volume | 25 | | | | | |

These PCR conditions have been optimized for our methods and preparation kits. Adoptions may be required.