


**EMMA ID: 05502**

**Gene: *Traf7***

**Common name: *TRAFF7 KO a conv***

### Allele Information



### Genotyping Information

Genotyping by end-point PCR based on gel is composed of a genespecific short range PCR using primers on wild type allele and a mutant allele-specific short range PCR. The combined results show the genotype of the mice. For example: mutant positive, wild type positive = Heterozygous.

#### PCR primer pairs and expected size bands

| Assay        | Forward Primer | Reverse Primer | Expected Size Band (bp) |
|--------------|----------------|----------------|-------------------------|
| control band | 1281_1         | 1281_2         | 335                     |
| Mutant       | 1535_32        | 1533_27        | 478                     |

#### Primer sequences

| Primer Name | Sequence 5' --> 3'        |
|-------------|---------------------------|
| 1535_32     | CTACTGACAGATTAATCATGGTAGC |
| 1533_27     | TGGGTCTCACACGGCATTCC      |
| 1281_1      | GTGGCACGGAACCTCTAGTC      |
| 1281_2      | CTTGTCAAGTAGCAGGAAGA      |

### PCR setup (Qiagen, Hot Start Plus)

| Component                     | Volume ( $\mu$ l) 1x | Final conc. |
|-------------------------------|----------------------|-------------|
| DNA (~ 50-100 ng)             | 2                    |             |
| Q-Solution (5x)               | 2,5                  | 0,5         |
| PCR-Buffer (10x)              | 2,5                  | 1           |
| DNTP mix (10 mM)              | 0,5                  | 0,2         |
| MgCl <sub>2</sub> (25 mM)     | 1,5                  | 1,5         |
| Primer 1 (10 pmol/ $\mu$ l)   | 1                    | 0,4         |
| Primer 2 (10 pmol/ $\mu$ l)   | 1                    | 0,4         |
| Taq Polymerase (5 U/ $\mu$ l) | 0,3                  | 0,06        |
| H <sub>2</sub> O*             | 13,7                 |             |
| Final volume                  | 25                   |             |


\* The amount of H<sub>2</sub>O is adjusted with the number of primer.

### Amplification conditions

| PCR Settings                                       | Temperature (°C)     | Time                       | # of cycles |
|--|----------------------|----------------------------|-------------|
| 1 Denaturation<br>(Melting)                        | 95°C                 | 5 min                      | 1           |
| 2 Amplification<br>(Melting, Annealing,<br>Polym.) | 94°C<br>60°C<br>72°C | 30 sec<br>45 sec<br>45 sec | 39          |
| 3 Polymerisation                                   | 72°C                 | 10 min                     | 1           |
| 4 Cooling  | 4°C                  | hold                       | 1           |

These PCR conditions have been optimized for our methods and preparation kits. Adoptions may be required.

### Gel Image



The fragment amplified with oligos 1 (1533\_27:  
**TGGGTCTCACACGGCATTCC**) + 5 (1535\_32:  
**CTACTGACAGATTAATCATGGTAGC**) detects  
heterozygous/homozygous conventional alleles.  
The wildtype fragment will not be detected  
applying the conditions described above.

Separated by gel electrophoresis on a 2% agarose gel.