



Genotyping protocol

Sept8 (IR00002287 / E23 ICS)

(ICS internal reference)

This report has been validated by:

Sylvie Jacquot, PhD, Head of Genotyping Service
33 (0)3 88 65 57 44
genotyping@igbmc.fr

The first version of this report was finalized the: 10 Oct 2010

For any question, please contact:

Institut Clinique de la Souris - ICS - Mouse Clinical Institute
1 rue Laurent Fries, BP 10142
67404 Illkirch Cedex, France
Email: genotyping@igbmc.fr
Web site: <http://www-mci.u-strasbg.fr/>

TABLE OF CONTENTS


Table of contents	2
1. Genotyping protocol and data	2
1.1. Genotyping strategy.....	2
1.2. PCR protocol.....	4
1.3. Picture of genotyping with various alleles.....	5
2. Cre and Flp genotyping method.....	Erreur ! Signet non défini.
2.1. Cre and Flp genotyping.....	Erreur ! Signet non défini.
2.2. PCR Protocol.....	Erreur ! Signet non défini.

1. Genotyping protocol and data

This section describes the condition used at the Mouse Clinical Institute (ICS) to genotype your **Sept8** Constitutive Knockout / Conditional Knockout (KO-cKO) project.

1.1. Genotyping strategy

The map below describes the position of the primers used for genotyping for each possible allele.



Sequence of primers used for genotyping:

Position	Primers	Sequence
Ef	3909	TGAGCCCCAGTTCTGGCTCCGT
Er	3910	AGCCAAGAGGCCAGGTTGGA
Kr	3277	CTCCTACATAGTTGGCAGTGTTGGG
L3f	3912	GCCTCTGCCAGGGTCAAGAATGGG
L3r	3911	GGGGATATGATGCCCTTCAGGCAGG
Lxr	3255	ACTGATGGCGAGCTCAGACCATAAC

PCR fragments expected size (bp):

Region analyzed	Primers used	Position on the primer <i>(see the map above)</i>	Targeted allele (KO) (L3)	cKO allele (L2)	KO allele (L-)	WildType allele (WT)
5' part of the selection marker	3909-3277	Ef / Kr	267	---	---	---
Presence of the distal loxP	3912-3911	L3f / L3r	233	233	---	255
Distal loxP specific PCR	3913-3255	L3f / Lxr	186	186	---	---
Excision of the selection marker	3909-3910	Ef / Er	5746*	346	---	225
Excision of the floxed exon(s), i.e. knock out	3909-3911	Ef / L3r	6670*	1270*	352**	1171**

*: this PCR product will not be observed using our PCR genotyping conditions (see description below)

**: this PCR is only verified if mice are generated

---: no Amplicon should be obtained

1.2. PCR protocol

This section describes the composition of the mix and cycling conditions used for genotyping.

Reagents:

- FastStart PCR Master (Roche)
- DNA (50ng/ μ l)
- 5' primer (100 μ M)
- 3' primer (100 μ M)
- Sterile H₂O

Volume:

- 7.5 μ l
- 1.5 μ l
- 0.06 μ l
- 0.06 μ l
- up to 15 μ l

Cycling conditions:


Temp	Time	#Cycles
95°C	4min	1
94°C	30s	
62°C	30s	34
72°C	1min	
72°C	7min	1
20°C	5min	1

NB: These PCR conditions have been optimized for high-throughput genotyping. Adaptation to small-scale may be required.

1.3. Picture of genotyping with various alleles

Analysis of PCR products pattern was not done by gel electrophoresis but using LabChip® 90 microfluidic apparatus. PCR products were run on the HT DNA 5K LabChip® 90 Assay Kit.

Representative genotyping picture



Note that as this technology is more sensitive than gel analysis, non specific signals and/or primer dimers may be visible on the picture.

2. Cre and Flp genotyping method

You will find the genotyping protocol in the publication:

[Highly-efficient, fluorescent, locus directed cre and FlpO deleter mice on a pure C57BL/6N genetic background.](#)

Birling MC, Dierich A, Jacquot S, Héault Y, Pavlovic G.
Genesis. 2012 Jun;50(6):482-9. doi: 10.1002/dvg.20826. Epub 2012 Mar 20.