



Genotyping protocol

1500015O10Rik

/ P5372

(ICS internal reference)

This report has been prepared by:

David Moulaert
genotyping@igbmc.fr

This report has been validated by:

Sylvie Jacquot, PhD, Head of Genotyping Service
33 (0)3 88 65 57 44
genotyping @igbmc.fr

The first version of this report was generated the: 03 September 2015

For any question, please contact:

Institut Clinique de la Souris - ICS - Mouse Clinical Institute
1 rue Laurent Fries, BP 10142
67404 Illkirch Cedex, France
Email: genotyping@igbmc.fr
Web site: <http://www-mci.u-strasbg.fr/>

TABLE OF CONTENTS


Table of contents	2
1. Genotyping protocol and data	2
1.1. Genotyping strategy.....	2
1.2. PCR protocol.....	4
2. Cre and Flp genotyping method.....	5

1. Genotyping protocol and data

This section describes the condition used at the Mouse Clinical Institute (ICS) to genotype your **1500015O10Rik** Constitutive Knockout / Conditional Knockout (KO-cKO x Cre) project.

1.1. Genotyping strategy

The map below describes the position of the primers used for genotyping for each possible allele.



Institut Clinique de la Souris
centre européen de recherche en biologie et en médecine
CERBgie

1 rue Laurent Fries - 67404 Illkirch Cedex - France - T. 33 (0)3 88 65 56 57 - F. 33 (0)3 88 65 56 90

Sequence of primers used for genotyping:

Position	Primers	Sequence
Ef	8383	CGGCTACAGCTCAGAACAAACCCAC
Er	8387	GCATTCTCTAGCCCCAGGCTATTAC
Kr	3209	CCAACAGCTTCCCCACAACCGG
L3f	8385	GGGCCTGAGATGGTAGACTTGCTTC
L3f ²	8384	TGAAGCGTGCAAACGACAGCT
L3r	8386	GGTTGTGAAGGTCAGTGGGTGTGC
Lxr	3255	ACTGATGGCGAGCTCAGACCATAAC
Rif	5966	GCACATGGCTGAATATCGACGGT

²: for a selected position, a second primer was designed

PCR fragments expected size (bp):

Region analyzed	Primers used	Position on the primer <i>(see the map above)</i>	Targeted allele (HR)	conditional allele (cKO)	KI allele	WildType allele
5' part of the selection marker	8383-3209	Ef / Kr	283	---	283	---
Presence of the distal loxP	8385-8386	L3f / L3r	393	393	---	378
Distal loxP specific PCR	8384-3255	L3f ² / Lxr	324	324	---	---
Excision of the selection marker	8383-8387	Ef / Er	7299*	395	---	232
Cre total excision	5966-3255	Ri1f / Lxr	3221*	---	471**	---

*: this PCR product will not be observed using our PCR genotyping conditions (see description below)

**: this PCR is only verified if mice are generated

---: no Amplicon should be obtained

1.2. PCR protocol

This section describes the composition of the mix and cycling conditions used for genotyping.

Reagents:

- FastStart PCR Master (Roche)
- DNA (50ng/ μ l)
- 5' primer (100 μ M)
- 3' primer (100 μ M)
- Sterile H₂O

Volume:

- 7.5 μ l
- 1.5 μ l
- 0.06 μ l
- 0.06 μ l
- up to 15 μ l

Cycling conditions:

Temp	Time	#Cycles
95°C	4min	1
94°C	30s	
62°C	30s	34
72°C	1min	
72°C	7min	1
20°C	5min	1

NB: These PCR conditions have been optimized for high-throughput genotyping. Adaptation to small-scale may be required.

2. Cre and Flp genotyping method

You will find the genotyping protocol in the publication:

[Highly-efficient, fluorescent, locus directed cre and FlpO deleter mice on a pure C57BL/6N genetic background.](#)

Birling MC, Dierich A, Jacquot S, Héault Y, Pavlovic G.
Genesis. 2012 Jun;50(6):482-9. doi: 10.1002/dvg.20826. Epub 2012 Mar 20.