



Genotyping protocol

Ttc39a

This report has been prepared by:

Christelle Roth
genotyping@igbmc.fr

This report has been validated by:

Sylvie Jacquot, PhD, Head of Genotyping Service
33 (0)3 88 65 57 44
genotyping @igbmc.fr

The first version of this report was generated the: 02 dec 2015

For any question, please contact:

Institut Clinique de la Souris - ICS - Mouse Clinical Institute
1 rue Laurent Fries, BP 10142
67404 Illkirch Cedex, France
Email: genotyping@igbmc.fr
Web site: <http://www-mci.u-strasbg.fr/>

TABLE OF CONTENTS


| | |
|----------------------------------------------|----------|
| Table of contents | 2 |
| 1. Genotyping protocol and data | 2 |
| 1.1. Genotyping strategy..... | 2 |
| 1.2. PCR protocol..... | 4 |
| 2. Cre and Flp genotyping method..... | 5 |

1. Genotyping protocol and data

This section describes the condition used at the Mouse Clinical Institute (ICS) to genotype your **Ttc39a** Constitutive Knockout / Conditional Knockout (KO-cKO x Cre) project.

1.1. Genotyping strategy

The map below describes the position of the primers used for genotyping for each possible allele.



Sequence of primers used for genotyping:

| Position | Primers | Sequence |
|----------|---------|------------------------------|
| Ef | 8610 | CCTGGGTGTTGTGTACTGATGGTGTGTC |
| Er | 8613 | GAGTCACACTCCCAACTCCCCCTGG |
| Kr | 3277 | CTCCTACATAGTTGGCAGTGTGGGG |
| L3f | 8611 | GACAAAGCCCGAGGTTCAGATTCC |
| L3r | 8612 | TCCAGGTGTTGCTGCATGAATCAAG |
| Lxr | 3255 | ACTGATGGCGAGCTCAGACCATAAC |
| Ri1f | 5966 | GCACATGGCTGAATATCGACGGT |

PCR fragments expected size (bp):

| Region analyzed | Primers used | Position on the primer <i>(see the map above)</i> | Targeted allele (HR) | conditional allele (cKO) | KI allele | WildType allele |
|----------------------------------|--------------|------------------------------------------------------|----------------------|--------------------------|-----------|-----------------|
| 5' part of the selection marker | 8610-3277 | Ef / Kr | 357 | --- | --- | --- |
| Presence of the distal loxP | 8611-8612 | L3f / L3r | 280 | 280 | --- | 200 |
| Distal loxP specific PCR | 8611-3255 | L3f / Lxr | 192 | 192 | --- | --- |
| Excision of the selection marker | 8610-8613 | Ef / Er | 7353* | 449 | --- | 323 |
| Cre total excision | 5966-3255 | Ri1f / Lxr | 3588* | --- | 471** | --- |

*: this PCR product will not be observed using our PCR genotyping conditions (see description below)

**: this PCR is only verified if mice are generated

---: no Amplicon should be obtained

1.2. PCR protocol

This section describes the composition of the mix and cycling conditions used for genotyping.

Reagents:

- FastStart PCR Master (Roche)
- DNA (50ng/ μ l)
- 5' primer (100 μ M)
- 3' primer (100 μ M)
- Sterile H₂O

Volume:

- 7.5 μ l
- 1.5 μ l
- 0.06 μ l
- 0.06 μ l
- up to 15 μ l

Cycling conditions:

| Temp | Time | #Cycles |
|------|------|---------|
| 95°C | 4min | 1 |
| 94°C | 30s | |
| 62°C | 30s | 34 |
| 72°C | 1min | |
| 72°C | 7min | 1 |
| 20°C | 5min | 1 |

NB: These PCR conditions have been optimized for high-throughput genotyping. Adaptation to small-scale may be required.

2. Cre and Flp genotyping method

You will find the genotyping protocol in the publication:

[Highly-efficient, fluorescent, locus directed cre and FlpO deleter mice on a pure C57BL/6N genetic background.](#)

Birling MC, Dierich A, Jacquot S, Héault Y, Pavlovic G.
Genesis. 2012 Jun;50(6):482-9. doi:10.1002/dvg.20826. Epub 2012 Mar 20.