

Name of Mouse model or mutation:

RBPJ-T223R-EM2-B6N

Description:

Point mutation model made using CRISPR/Cas9.

Type of mutation:


SNP: T223R

Sequence details**Rbpj WT**

```
AGGCTAAATGTTTGCCACCAGAATCTGTTTGTTATTTGCATTACTGTAAAAAGTATATATAATTTG  
CCAAGCCAAAGCCCCTTTCTTTGTGCGTGCCTCCCCGCATCTAAATAGGATCCCCCTGTCTTTCAGTGT  
GCATTGCTTCAGGAACGAAGGTGGCACTGTTCAATCGCCTTCGGTCCCAGACAGTTAGTACCAGGTA  
CCTGCATGTAGAAGGAGGGAATTTCCACGCCAGTTCACAACAGTGGGGAGCATTTTACATCCATCTC  
TGTGAGTACAAAATGTACATTTGGTATTTTTAGTGTGAAAATAATATGTGCTCCTTCTTGGGTTAATA
```

RBPJ-T223R-EM2-B6N

```
AGGCTAAATGTTTGCCACCAGAATCTGTTTGTTATTTGCATTACTGTAAAAAGTATATATAATTTG  
CCAAGCCAAAGCCCCTTTCTTTGTGCGTGCCTCCCCGCATCTAAATAGGATCCCCCTGTCTTTCAGTGT  
GCATTGCTTCAGGAACGAAGGTGGCACTGTTCAATCGCCTTCGGTCCCAGAGAGTAAAGTACCAGGT  
ACCTGCATGTAGAAGGAGGGAATTTCCACGCCAGTTCACAACAGTGGGGAGCATTTTACATCCATCT  
CTGTGAGTACAAAATGTACATTTGGTATTTTTAGTGTGAAAATAATATGTGCTCCTTCTTGGGTTAAT  
A
```

RBPJ-T223R-EM2-B6N Heterozygous F1 animal sequence trace:

Nucleotide Alignment:

```

                *      20      *      40      *      60      *      80      *      100     *      120
Rbpj_WT      : AGGCTAAAATGTTTGCACCCAGAATCTGTTTGTATTGTCATTACTGTAAAAAGTATATATAATTTGCCAAGCCAAAGCCCCTTCTTTGTGCGTGCCTCCCCGCATCTAAATAGGATCCCCCTG
Rbpj_T223R  : AGGCTAAAATGTTTGCACCCAGAATCTGTTTGTATTGTCATTACTGTAAAAAGTATATATAATTTGCCAAGCCAAAGCCCCTTCTTTGTGCGTGCCTCCCCGCATCTAAATAGGATCCCCCTG
                *      140     *      160     *      180     *      200     *      220     *      240     *
Rbpj_WT      : TCTTTCAGTGTGCATTGCTTCAGGAACGAAGGTGGCACTGTTCAATCGCCTTCGGTCCCAGACAGTTCAGTACCAGGTACCTGCATGTAGAAGGAGGGAATTTCCACGCCAGTTCACAACAGTGGGG
Rbpj_T223R  : TCTTTCAGTGTGCATTGCTTCAGGAACGAAGGTGGCACTGTTCAATCGCCTTCGGTCCCAGACAGTTCAGTACCAGGTACCTGCATGTAGAAGGAGGGAATTTCCACGCCAGTTCACAACAGTGGGG
                *      260     *      280     *      300     *      320     *
Rbpj_WT      : AGCATTTTACATCCATCTCTGTGAGTACAAAATGTACATTTGGTATTTTGTAGTGTGAAAATAATATGTGCTCCTTCTTGGGTTAATA : 339
Rbpj_T223R  : AGCATTTTACATCCATCTCTGTGAGTACAAAATGTACATTTGGTATTTTGTAGTGTGAAAATAATATGTGCTCCTTCTTGGGTTAATA : 339
                AGCATTTTACATCCATCTCTGTGAGTACAAAATGTACATTTGGTATTTTGTAGTGTGAAAATAATATGTGCTCCTTCTTGGGTTAATA

```

Predicted Protein Alignment:

```

                *      20      *      40
Rbpj_WT      : VCIASGTKVALFNRLRSQVSTRYLHVEGGNFHASSQQWGAIFYIHL
Rbpj_T223R  : VCIASGTKVALFNRLRSQVSTRYLHVEGGNFHASSQQWGAIFYIHL
                VCIASGTKVALFNRLRSQ VSTRYLHVEGGNFHASSQQWGAIFYIHL

```

QC strategy employed at Harwell to check the edited allele:

Genomic DNA was extracted from ear clip biopsies and amplified in a PCR reaction using the following conditions/primer sequences:

Geno_Rbpj_F1	AGGCTAAAATGTTTGCCACCAG
Geno_Rbpj_R1	TATTAACCCAAGAAGGAGCACA
Taq Polymerase used	Roche Expand Long Range DNTPack
Annealing Temperature (°C)	56
Elongation time (min)	0.5
WT product size (bp)	339
Mutant product size (bp)	339
Notes	Three off-target sites for the sgRNA used were also checked and confirmed as WT. The primers are detailed in table below.

OT 1 - Rbpsuh rs3	
Geno_Rbpsuh_F1	GCACAGAAGTCTTACGGAAATGAAA
Geno_Rbpsuh_R1	GGGCCCATCCCTCATAGAAC
Annealing Temperature (°C)	62
Elongation Time (min)	1
WT size (bp)	872
Mut size (bp)	872
OT 2 - Rpbsuh ps1	
Geno_Rbpsuh_Ps1_F1	TATCAATCTTGGGAGCGCCA
Geno_Rbpsuh_Ps1_R1	TGTTCTCAACAAACCCTGTGAGT
Annealing Temperature (°C)	62
Elongation Time (min)	1
WT size (bp)	508
Mut size (bp)	508
OT 3 - Rbpsuh ps2	
Geno_Rbpsuh_Ps2_F1	CCCGGCGTTTCAGCCTAAT
Geno_Rbpsuh_Ps2_R1	TGAACAAGAGTCTCAACCCTGTG
Annealing Temperature (°C)	62
Elongation Time (min)	1
WT size (bp)	540
Mut size (bp)	540

All amplicons were sent for Sanger sequencing to check for integration of the donor oligo sequence at the target site. F1 sequences should be heterozygous unless on Y chromosome.



Copy counting of the donor sequence was carried out by ddPCR at the F1 stage to confirm donor oligos were inserted once on target into the genome. The following Taqman assay was used to copy count the donor sequence compared against a VIC-labelled reference assay for Dot1l:

Assay name	Rbpj-T223R-Donor4-MUT1
Forward Primer	GGAACGAAGGTGGCACTGTT
Reverse Primer	CATGCAGGTACCTGGTACTTACTC
Probe	CAATCGCCTTCGGTCCCAGA
Label	FAM-BHQ1

The ddPCR assay is specific to the T223R mutation of RBPJ and the WT gene is not recognised by this assay. WT controls are expected to call at 0 copies and a single integration for a correct mutation is expected to call at 1 copy for F1 (HET) animals.


Rbpj-T223R Genotyping Strategy

Animals have been engineered using the CRISPR/Cas9 technology. Most of the knockout alleles generated through this method will be obtained by deletion of a critical exon or by introduction of an indel (insertion/deletion) within the coding sequence of a critical exon (see picture below).




qPCR genotyping strategy

Standard PCR is the amplification of DNA between a pair of primers. Quantitative PCR employs the same principal as standard PCR, although it actually monitors the progress of the DNA synthesis as it occurs. The progress of the reaction is measured by using a Taqman probe. This is a short DNA oligo that is complimentary to part of the DNA sequence between the forward and reverse primers. At the 5' end of the probe there is a fluorescent reporter (R) and at the 3' end a quencher (Q). Whilst they are in close contact with each other there is no fluorescent signal.



As the forward primer is extended the reporter is cleaved from the probe resulting in a fluorescent signal being detected. Once the primer extends enough to release the quencher this signal is blocked. By using probes with different fluorescent signals multiple PCR assays can be multiplexed and run together.



Allele specific primer and probe amplification assay (ASPPAA) PCR

This is a new real-time PCR method (Billard *et al.*, 2012) in which an allele specific primer and an allele specific probe designed specific to the SNPs. The primer is designed such a way that its 3' end ends with a specific SNP. The probe is also designed specific to the SNPs at its 3' end giving a primer probe overlap. A maximum of 3nt overlap between a primer and probe is allowed.





Figure1: Figure showing principle of ASPPAA PCR. The dotted line indicates the position of the SNP.

PCR reaction plot



The number of cycles the PCR takes to reach a set threshold is known as the CT value. This is inversely correlated to the amount of template DNA in the sample.

e.g. CT 25 = 2 x template DNA
CT 26 = 1 x template DNA
CT above 30 = no template represented in the sample

CT value can be used to determine how many copies of a particular allele samples have.

All our qPCR are run in duplicate. A FAM labelled genotyping assay is run in multiplex with a VIC labelled internal control Dot1l.

References:

Billard A., Laval V., Fillinger S., Leroux P., Lachaise H., Beffa R., et al. (2012).
The allele-specific probe and primer amplification assay, a new real-time PCR method for fine quantification of single-nucleotide polymorphisms in pooled DNA.
Appl. Environ. Microbiol. 78 1063–1068. 10.1128/AEM.06957-11



Rbpj-T223R Genotyping Strategy

Samples are genotyped with both WT and Mutant assays. These are FAM labelled assays that are designed to detect the critical exon that has been modified. If the animal contains the modified allele the copy number of the WT assay should drop by 1 and the mutant assay should raise by 1. For autosomal genes that have been targeted this means the following

- WT= 2 copies of the WT assay and 0 copies of the Mutant assay
- HET = 1 copy of the WT assay and 1 copy of the Mutant assay
- HOM = 0 copies of the WT assay and 2 copies of the Mutant assay

Fragment Sequences

The highlighted SNPs below are the CRISPR/Cas9 modifications for the Rbpj gene.

Rbpj WT CCCAGACAGTTAGTACCAGGTA
 Rbpj-T223R CCCAGAGAGTAAGTACCAGGTA

Rbpj-T223R-WT2 assay (FAM labelled probe)

CCAAGCCAAAGCCCCTTTCTTTGTGCGTGCCCTCCCCGCATCTAAATAGGATCCCCCTGTCTTTTCAGTG
 TGCATTGCTTCAGGAACGAAGGTGGCACTGTTCAATCGCCTTCGGTCCCAGACAGTTAGTACCAGGTA
CCTGCCATGTAGAAGGAGGGAATTTCCACGCAGTTTACAACAGTGGGGAGCATTTTACATCCATCTCT
 GTGAGTACAAAATGTACATTTGGTATTTTTAGTGTGAAAATAATATGTGCTCCTTCTTGGGTTAATA

Rbpj-T223R-LOA-WT1 primers and probe

Primer 1 = GCGTGGAATTCCTCCTTCTAC
 Primer 2 = AATCGCCTTCGGTCCCAGAC
 Probe = CAGGTACCTGGTACTAACTGTC } Allele specific primer and probes

Rbpj-T223R-Donor4-MUT1 assay (FAM labelled probe)

CCAAGCCAAAGCCCCTTTCTTTGTGCGTGCCCTCCCCGCATCTAAATAGGATCCCCCTGTCTTTTCAGTG
 TGCATTGCTTCAGGAACGAAGGTGGCACTGTTCAATCGCCTTCGGTCCCAGAGTAAAGTACCAGGTA
CCTGCCATGTAGAAGGAGGGAATTTCCACGCCAGTTTACAACAGTGGGGAGCATTTTACATCCATCTCT
 GTGAGTACAAAATGTACATTTGGTATTTTTAGTGTGAAAATAATATGTGCTCCTTCTTGGGTTAATA

Rbpj-T223R-LOA-WT1 primers and probe

Primer 1 = GGAACGAAGGTGGCACTGTT
 Primer 2 = CATGCAGGTACCTGGTACTTACTC
 Probe = CAATCGCCTTCGGTCCAGA

Dot1l internal control (VIC labelled)

TCATAGGGTGACTGGCCAACCCAGGGAAGCCGGAGTGCTGCGTCTTCTGTTTCCCTTGTCTTTTCCCCTCTAGTC
 GTTTTCTGTTAGTAGTTGGCATCCTTATGCTTCATCTTACAGTCGACTTGAGAGCTGGCCCTGAATGGTCGTGCT
GGGGCAAGGCTTTATTTTCAGGCGTAGCACACATGGTGGCCAATGGGACTCTGTAGGATCTGCCACACCCATCAG


Primer 1 = GCCCAGCACGACCATT
 Primer 2 = TAGTTGGCATCCTTATGCTTCATC
 Probe = CCAGCTCTCAAGTCG



qPCR master mix

ABI GTX Taqman master mix	5µl
Primers Dot1L_2F (20µM)	0.225µl
Primers Dot1L_R (20µM)	0.225µl
Probe DotL_2M (5µM)	0.2µl
FAM Assay (probe 5µM & primers 15µM each)	0.3µl
ddH2O	1.55µl
DNA (1/10 dilution of ABI Sample-to-SNP prep)	2.5µl

Rbpj-T223R Mutant copy called result, image showing both replicates and controls



Version No. 1
Date: 10/01/19
Created/Updated by: Ramakrishna Kurapati
Approved by: Daniel Ford